

LISTING OF THE CLAIMS:


1. – 19. (Canceled)

20. (Withdrawn) A method according to Claim 17 wherein the CB1 receptor antagonist is N-piperidino-5 (4-bromophenyl)-1 – (2,4-dichlorophenyl) -4-ethylpyrazole-4-ethylpyrazole-3-carboxamide or one of its pharmaceutically acceptable salt.

21. – 27. (Canceled)

28. (New) A method of treating hepatic fibrosis in a mammal in need thereof which comprises administering a therapeutically effective amount of at least one CB1 receptor antagonist.

29. (New) The method of Claim 28 wherein the CB1 receptor is a compound of Formula II:



Formula II

or a pharmaceutically acceptable salt thereof, wherein

g_2, g_3, g_4, g_5 and g_5 and w_2, w_3, w_4, w_5 and w_6 are identical or different and are independently hydrogen, a chlorine or bromine atom, a (C_1-C_3) alkyl, a (C_1-C_3) alkoxy, a trifluoromethyl or a nitro group and g_4 is optionally a phenyl group;

R₄ is hydrogen or a (C₁-C₃) alkyl;

X is either a direct bond or a group $-(CH_2)_xN(R_3)-$ in which R₃ is hydrogen or a (C₁-C₃) alkyl and x is zero or one;

R is a group-NR₁R₂ in which R₁ and R₂ are independently a (C₁-C₆)-alkyl; an non-aromatic (C₃-C₁₅) carbocyclic radical which is optionally substituted, said substituent (s) being other than a substituted carbonyl; an amino (C₁-C₄) alkyl group in which the amino is optionally disubstituted by a (C₁-C₃) alkyl; a cycloalkyl (C₁-C₃) alkyl group in which the cycloalkyl is C₃-C₁₂; a phenyl which is unsubstituted or monosubstituted or polysubstituted by a halogen, by a (C₁-C₅) alkyl or by a (C₁-C₅) alkoxy; a phenyl (C₁-C₃) alkyl; a diphenyl (C₁-C₃) alkyl; a naphthyl; an anthracenyl; a saturated 5- to 8-membered heterocyclic radical which in unsubstituted or substituted by a (C₁-C₃) alkyl; by a hydroxyl or by a benzyl; a 1-adamantylmethyl; an aromatic heterocycle which is unsubstituted or monosubstituted or polysubstituted by a halogen, by a (C₁-C₅) alkyl or by a (C₁-C₅) alkoxy; a (C₁-C₅) alkyl which is substituted by an aromatic heterocycle which is unsubstituted or monosubstituted or polysubstituted by a halogen, by a (C₁-C₅) alkyl or by a (C₁-C₅) alkoxy; or else R is hydrogen and R₂ is as defined above; or else R₁ and R₂ form a saturated 5- to 8-membered heterocyclic radical with the nitrogen atom to which they are bonded, said heterocyclic radical being other than morpholine when w₂, w₃, w₄, w₅, w₆, g₂, g₃, g₄, g₅ and g₆ are all hydrogen; a group R₂ as defined above when X is $-(CH_2)_xN(R_3)-$; a group R₅ when X is a direct bond, R₅ being a (C₁-C₃) alkyl; a (C₃-C₁₂) cycloalkyl which is unsubstituted or substituted by a (C₁-C₅) alkyl; a phenyl (C₁-C₃) alkyl which is unsubstituted or substituted by a halogen or by a (C₁-C₅) alkyl;

a cycloalkyl (C₁-C₃) alkyl in which the cycloalkyl is C₃-C₁₂ and is unsubstituted by a (C₁-C₅) alkyl; or a 2-norbornylmethyl.

30. (New) The method according to Claim 28 wherein the CB1 receptor antagonist is N-piperidino-3-pyrazolecarboxamide or a pharmaceutically acceptable salt thereof.

31. (New) The method of Claim 28 wherein the CB1 receptor antagonist is N-piperidino-5-(4-bromophenyl)-1-(2,4-dichlorophenyl)-4-ethylpyrazole-3-carboxamide or a pharmaceutically acceptable salt thereof.

32. (New) The method of Claim 28 wherein the CB1 receptor antagonist is N-piperidino-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methylpyrazole-3-carboxamide or a pharmaceutically acceptable salt thereof.

33. (New) The method of Claim 28 wherein the daily dosage of CB1 receptor antagonist is from 0.01 mg to 500 mg.

34. (New) The method of Claim 33 wherein the daily dosage of CB1 receptor antagonist is from 1 mg to 100 mg.

35. (New) The method of Claim 28 wherein the CB1 receptor is selected from the group consisting of:

- a) a protein having an amino acid sequence comprising SEQ ID NO: 1 or a portion of SEQ ID NO: 1, having the biological function of a G protein-coupled cellular receptor, capable of binding THC and transducing a cellular signal;
- b) a protein having an amino acid sequence comprising SEQ ID NO: 2 or a portion of SEQ ID NO: 2, having the biological function of a G protein-coupled cellular receptor, capable of binding THC and transducing a cellular signal;
- c) an allele of the protein having the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 2, having the biological function of a G protein-coupled cellular receptor, capable of binding THC and transducing a cellular signal;
- d) a protein having the amino acid sequence of SEQ ID NO: 1 with a Phenylalanine to Leucine substitution at position 200; and/or Isoleucine to Valine substitution at position 216; and/or a Valine to Alanine substitution at position 246;
- e) a protein having the amino acid sequence of SEQ ID NO: 2 with a Phenylalanine to Leucine substitution at position 139; and/or an Isoleucine to Valine substitution at position 155; and/or a Valine to Alanine substitution at position 185; and
- f) a protein comprising the amino acid sequences of SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, and SEQ ID NO: 9 or amino acid sequences 80% homologous to these, said protein having the biological function of a G protein-coupled cellular receptor, capable of binding THC and transducing a cellular signal.

36. (New) The method of Claim 28 wherein the CB1 receptor is a protein having a homology at the amino acid level with SEQ ID NO: 1 of at least 45% having the biological

function of a G protein-coupled cellular receptor, capable of binding THC and transducing a cellular signal.

37. (New) The method of Claim 36 wherein the homology is at least 60%.